

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1600RXA

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 May 10 PROUSDDR now available on STN  
NEWS 4 May 19 PROUSDDR: One FREE connect hour, per account, in both May  
and June 2004  
NEWS 5 May 12 EXTEND option available in structure searching  
NEWS 6 May 12 Polymer links for the POLYLINK command completed in REGISTRY  
NEWS 7 May 17 FRFULL now available on STN  
NEWS 8 May 27 New UPM (Update Code Maximum) field for more efficient patent  
SDIs in CApplus  
NEWS 9 May 27 CApplus super roles and document types searchable in REGISTRY  
NEWS 10 May 27 Explore APOLLIT with free connect time in June 2004  
NEWS 11 Jun 22 STN Patent Forums to be held July 19-22, 2004  
  
NEWS EXPRESS MARCH 31 CURRENT WINDOWS VERSION IS V7.00A, CURRENT  
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 26 APRIL 2004  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 12:55:16 ON 23 JUN 2004

=> fil reg

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 12:55:42 ON 23 JUN 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 22 JUN 2004 HIGHEST RN 697737-72-7  
DICTIONARY FILE UPDATES: 22 JUN 2004 HIGHEST RN 697737-72-7

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 6, 2004


Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\STNEXP4\QUERIES\10637190.str



chain nodes :

7 8 15 16

ring nodes :

1 2 3 4 5 6 9 10 11 12 13 14 17 19 20 21 22 23 24 25 26 27 28

chain bonds :

4-7 7-8 8-9 12-15 15-16 16-17 22-23

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 9-10 9-14 10-11 11-12 12-13 13-14 17-19 17-22  
19-20 20-21 21-22 23-24 23-28 24-25 25-26 26-27 27-28

exact/norm bonds :

1-2 1-6 2-3 3-4 4-5 4-7 5-6 7-8 8-9 9-10 9-14 10-11 11-12 12-13 12-15  
13-14 15-16 16-17 17-19 17-22 19-20 20-21 21-22 22-23 23-24 23-28 24-25  
25-26 26-27 27-28

G1:O,N

G2:C,N

Match level :


1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:Atom 10:Atom  
11:Atom 12:Atom 13:Atom 14:Atom 15:CLASS 16:CLASS 17:Atom 19:Atom 20:Atom  
21:Atom 22:Atom 23:Atom 24:Atom 25:Atom 26:Atom 27:Atom 28:Atom

L1 STRUCTURE UPLOADED

=> d

L1 HAS NO ANSWERS

L1 STR



G1 O,N

G2 C,N

Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 13:00:07 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 1 TO ITERATE

100.0% PROCESSED 1 ITERATIONS

1 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*

BATCH \*\*COMPLETE\*\*

PROJECTED ITERATIONS: 1 TO 80

PROJECTED ANSWERS: 1 TO 80

L2 1 SEA SSS SAM L1

=> s l1 full

FULL SEARCH INITIATED 13:00:10 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 21 TO ITERATE

100.0% PROCESSED 21 ITERATIONS

15 ANSWERS

SEARCH TIME: 00.00.01

L3 15 SEA SSS FUL L1

=> s l3 and caplus/lc


35942085 CAPLUS/LC

L4 10 L3 AND CAPLUS/LC


=> s 13 not 14  
L5 5 L3 NOT L4

=> d 15 1-5

L5 ANSWER 1 OF 5 REGISTRY COPYRIGHT 2004 ACS on STN  
 RN 463316-13-4 REGISTRY  
 CN 4-Isokazolecarboxylic acid, 3-(2-chlorophenyl)-5-methyl-,  
 4-[2-(2,4-dinitrophenyl)ethenyl]-2-methoxyphenyl ester (9CI) (CA INDEX  
 NAME)  
 FS 3D CONCORD  
 MF C26 H18 Cl N3 O8  
 SR Chemical Library  
 LC STN Files: CHEMCATS




PAGE 1-A



PAGE 2-A


\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L5 ANSWER 2 OF 5 REGISTRY COPYRIGHT 2004 ACS on STN  
 RN 462624-31-3 REGISTRY  
 CN 4-Isokazolecarboxylic acid, 3-(2-chlorophenyl)-5-methyl-,  
 2-methoxy-4-(2-phenylethenyl)phenyl ester (9CI) (CA INDEX NAME)  
 FS 3D CONCORD  
 MF C26 H20 Cl N O4  
 SR Chemical Library  
 LC STN Files: CHEMCATS




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L5 ANSWER 3 OF 5 REGISTRY COPYRIGHT 2004 ACS on STN  
 RN 462611-49-0 REGISTRY  
 CN 4-Isokazolecarboxylic acid, 3-(2-chlorophenyl)-5-methyl-,  
 4-(2-cyano-2-phenylethenyl)-2-methoxyphenyl ester (9CI) (CA INDEX NAME)  
 FS 3D CONCORD  
 MF C27 H19 Cl N2 O4  
 SR Chemical Library  
 LC STN Files: CHEMCATS



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L5 ANSWER 4 OF 5 REGISTRY COPYRIGHT 2004 ACS on STN  
 RN 302574-97-6 REGISTRY  
 CN 4-Isokazolecarboxylic acid, 3-(2-chlorophenyl)-5-methyl-,  
 4-[2-(4-chlorophenyl)-2-cyanoethenyl]-2-methoxyphenyl ester (9CI) (CA  
 INDEX NAME)  
 FS 3D CONCORD  
 MF C27 H18 Cl2 N2 O4  
 SR Chemical Library  
 LC STN Files: CHEMCATS



PAGE 1-A




PAGE 2-A

\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

LS ANSWER 5 OF 5 REGISTRY COPYRIGHT 2004 ACS on STN  
RN 219865-50-6 REGISTRY  
CN 4-isoxazolecarboxylic acid, 3-(2,6-dichlorophenyl)-5-methyl-,  
4-[1-cyano-2-[4-(trifluoromethyl)phenyl]ethenyl]phenyl ester (9CI) (CA  
INDEX NAME)  
FS 3D CONCORD  
MF C27 H15 Cl2 F3 N2 O3  
SR CAS Client Services  
LC STN Files: CHEMCATS

PAGE 1-A



PAGE 2-A



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

=> fil caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

172.06

172.27

FILE 'CAPLUS' ENTERED AT 13:01:18 ON 23 JUN 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 23 Jun 2004 VOL 140 ISS 26

FILE LAST UPDATED: 22 Jun 2004 (20040622/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d his

(FILE 'HOME' ENTERED AT 12:55:16 ON 23 JUN 2004)

FILE 'REGISTRY' ENTERED AT 12:55:42 ON 23 JUN 2004

L1 STRUCTURE UPLOADED

L2 1 S L1

L3 15 S L1 FULL

L4 10 S L3 AND CAPLUS/LC

L5 5 S L3 NOT L4

FILE 'CAPLUS' ENTERED AT 13:01:18 ON 23 JUN 2004

=> s l4

L6 19 L4

=> s l4 and Farnesoid

19 L4

253 FARNESOID

3 FARNESOIDS

255 FARNESOID

(FARNESOID OR FARNESOIDS)

L7 16 L4 AND FARNESOID

=> d ibib abs hitstr 1016

16 ANSWERS ARE AVAILABLE. SPECIFIED ANSWER NUMBER EXCEEDS ANSWER SET SIZE

The answer numbers requested are not in the answer set.

ENTER ANSWER NUMBER OR RANGE (1):1-16





L7 ANSWER 1 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 2004:453343 CAPLUS  
TITLE: Crystal structure of the human **farnesoid X** receptor ligand binding domain complexed with fexaramine and identification and development of novel small molecule ligands for FXR  
INVENTOR(S): Downes, Michael R.; Verdicia, Mark A.; Noel, Joseph P.; Evans, Ronald M.; Bowman, Lindsey J.; Bowman, Marianne  
PATENT ASSIGNEE(S): The Salk Institute for Biological Studies, USA  
SOURCE: PCT Int. Appl., 139 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

| PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE     |
|---|------|----------|-----------------|----------|
| WO 2004046323   | A2   | 20040603 | WO 2003-US36548 | 20031114 |
| V: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ |      |          |                 |          |
| RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG  |      |          |                 |          |

PRIORITY APPLN. INFO.: US 2002-426665P P 20021115  
US 2002-426668P P 20021115


AB The present invention provides compns. comprising the ligand binding domain (LBD) of a human **farnesoid X** receptor (FXR) in crystalline form. In alternative embodiments, the LBD of FXR is complexed with a ligand therefor. There are provided high resolution structures and structure coordinates of FXR complexed with a novel high affinity agonist, fexaramine. The discovered structure of a FXR LBD provides the first three-dimensional view of the structural basis for FXR ligand binding. The present invention further provides a computer for producing a three-dimensional representation of FXR or a complex thereof, and a computer for determining at least a portion of the structure coordinates of FXR or a complex thereof. The present invention further provides methods of using this structural information to predict mol. capable of binding to FXR; to identify compds. with agonist, antagonist or partial agonist activity for FXR; and to determine whether a test compound is capable of binding to the LBD of FXR. The present invention further provides compns. comprising compds. identified by such invention methods. Identification and development of novel small mol. ligands for FXR, and activation of FXR and induction of FXR target genes by these novel compds. is disclosed.

IT INDEXING IN PROGRESS  
IT 278779-30-9E, GW4064  
RL: BSU (Biological study, unclassified); CPN (Combinatorial preparation);

L7 ANSWER 1 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)  
THU (Therapeutic use); BIOL (Biological study); CMBI (Combinatorial study); PREP (Preparation); USES (Uses)  
(FXR ligand; crystal structure of human **farnesoid X** receptor ligand binding domain complexed with fexaramine and identification and development of novel small mol. ligands for FXR)

RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A

HO<sub>2</sub>C

L7 ANSWER 2 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 2004:453231 CAPLUS  
TITLE: Non-steroidal fkr agonists  
INVENTOR(S): Nicolaou, Kyriacos C.; Roecker, Anthony J.; Hughes, Robert; Pfefferkorn, Jeffrey A.  
PATENT ASSIGNEE(S): The Scripps Research Institute, USA  
SOURCE: PCT Int. Appl., 75 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

| PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE     |
|---|------|----------|-----------------|----------|
| WO 2004046162   | A2   | 20040603 | WO 2003-US36195 | 20031114 |
| V: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD |      |          |                 |          |
| RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG  |      |          |                 |          |


PRIORITY APPLN. INFO.: US 2002-426456P P 20021114  
US 2003-491185P P 20030729

AB Abstract Potent non-steroidal **farnesoid X** receptor (FXR) agonists are N-aryl-N-arylmethyl amide and ureido compds. having the chemical structure represented by the following formula (I): INSERT FORMULA where in E1 is (C1-C8)alkyl, cyclohexyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, Ph, or NH(C1-C8)alkyl; L1 and L2 are both H, or together form a pi-bond; X1 is C(O), or CH<sub>2</sub>; Y1 is H, NH<sub>2</sub>, NH(ZZ)23, or OZ<sub>4</sub>; aryl moiety A1 is selected from the group of radicals consisting of: INSERT FORMULA A2 and G1 - G11 are as defined in the specification; and T1 and T2 are each independently O, S, NH, or N(C1-C8)alkyl. The FXR agonists are useful as therapeutic agents for the treatment of diseases linked to cholesterol, bile acids, and the regulation of metabolism and homeostasis.

IT INDEXING IN PROGRESS  
IT 278779-30-9, GW 4064  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(non-steroidal fkr agonists)

RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A

HO<sub>2</sub>C

L7 ANSWER 3 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 2003:973413 CAPLUS  
DOCUMENT NUMBER: 140:229012  
TITLE: Hepatoprotection by the farnesoid X receptor agonist GW4064 in rat models of intra- and extrahepatic cholestasis  
AUTHOR(S): Liu, Yaping; Binz, Jane; Numerick, Mary Jo; Dennis, Steve; Luo, Guizhen; Desai, Bhasha; MacKenzie, Kathleen I.; Mansfield, Traci A.; Kliever, Steven A.; Goodwin, Bryan; Jones, Stacey A.  
CORPORATE SOURCE: Nuclear Receptor Functional Analysis, High Throughput Biology, GlaxoSmithKline, Research Triangle Park, NC, USA  
SOURCE: Journal of Clinical Investigation (2003), 112(11), 1678-1687  
CODEN: JCINAO; ISSN: 0021-9738  
PUBLISHER: American Society for Clinical Investigation  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB **Farnesoid X receptor (FXR)** is a bile acid-activated transcription factor that is a member of the nuclear hormone receptor superfamily. FXR-null mice exhibit a phenotype similar to Byler disease, an inherited cholestatic liver disorder. In the liver, activation of FXR induces transcription of transporter genes involved in promoting bile acid clearance and represses genes involved in bile acid biosynthesis. We investigated whether the synthetic FXR agonist GW4064 could protect against cholestatic liver damage in rat models of extrahepatic and intrahepatic cholestasis. In the bile duct-ligation and  $\alpha$ -naphthylisothiocyanate models of cholestasis, GW4064 treatment resulted in significant reductions in serum alanine aminotransferase, aspartate aminotransferase, and lactate dehydrogenase, as well as other markers of liver damage. Rats that received GW4064 treatment also had decreased incidence and extent of necrosis, decreased inflammatory cell infiltration, and decreased bile duct proliferation. Anal. of gene expression in livers from GW4064-treated cholestatic rats revealed decreased expression of bile acid biosynthetic genes and increased expression of genes involved in bile acid transport, including the phospholipid flippase MDR2. The hepatoprotection seen in these animal models by the synthetic FXR agonist suggests FXR agonists may be useful in the treatment of cholestatic liver disease.  
IT 278779-30-9, GW4064  
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(hepatoprotection by farnesoid X receptor agonist GW4064 in rat models of intra- and extrahepatic cholestasis)  
RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 3 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A

REFERENCE COUNT: 53 THERE ARE 53 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 4 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 2003:855658 CAPLUS  
DOCUMENT NUMBER: 139:317457  
TITLE: Compositions and methods using farnesoid X receptor ligands for hepatoprotection and treatment of cholestasis  
INVENTOR(S): Kliever, Steven Anthony; Willson, Timothy Mark  
PATENT ASSIGNEE(S): USA  
SOURCE: U.S. Pat. Appl. Publ., 8 pp.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


| PATENT NO.    | KIND | DATE     | APPLICATION NO. | DATE     |
|---------------|------|----------|-----------------|----------|
| US 2003203939 | A1   | 20031030 | US 2002-132311  | 20020425 |
| WO 2003090745 | A1   | 20031106 | WO 2003-US10519 | 20030407 |

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OH, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM  
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, EE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2002-132311 A 20020425  
OTHER SOURCE(S): MARPAT 139:317457  
AB Methods for the treatment of cholestatic liver disease and reduction and prevention of hepatic injury resulting from cholestasis via administration of a FXR ligand are provided. Bile duct-ligated rats treated with FXR ligand GW4064 had a pronounced improvement in liver function as defined by a reduction in a panel of liver disease serum marker enzymes.  
IT 278779-30-9, GW4064  
RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(FXR agonist; farnesoid X receptor ligands for hepatoprotection and treatment of cholestasis)  
RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 4 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A


L7 ANSWER 5 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN  
 ACCESSION NUMBER: 2003:777952 CAPLUS  
 DOCUMENT NUMBER: 139:286360  
 TITLE: Methods using **farnesoid X receptor (FXR)** agonists for weight loss and alteration of cell metabolism  
 INVENTOR(S): Jones, Stacey Ann; Klierer, Steven Anthony; Mansfield, Traci Ann  
 PATENT ASSIGNEE(S): Smithkline Beecham Corporation, USA; Curagen Corporation  
 SOURCE: PCT Int. Appl., 25 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

| PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE     |
|---|------|----------|-----------------|----------|
| WO 2003080803   | A2   | 20031002 | WO 2003-US8634  | 20030319 |
| W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GE, GD, GR, GM, GU, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, OM, PE, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM<br>RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG |      |          |                 |          |

PRIORITY APPLN. INFO.: MARPAT 139:286360  
 OTHER SOURCE(S):  
 AB Treatment of human hepatocytes with **farnesoid X receptor (FXR)** agonists resulted in increased expression of FGF-19. Methods of using FXR agonists to alter cell metabolism, and in pharmaceutical weight loss methods, are described.  
 IT 278779-30-9, GW4064 278779-30-9D, GW4064, amino acid conjugates  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (farnesoid X receptor agonists for weight loss and alteration of cell metabolism)  
 RN 278779-30-9 CAPLUS  
 CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 5 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A




PAGE 2-A

RN 278779-30-9 CAPLUS  
 CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 5 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A




L7 ANSWER 6 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:723027 CAPLUS  
 DOCUMENT NUMBER: 139:286515  
 TITLE: Estrogen receptor  $\alpha$  regulates expression of the orphan receptor small heterodimer partner  
 AUTHOR(S): Lai, KehDi; Harnish, Douglas C.; Evans, Mark J.  
 CORPORATE SOURCE: Wyeth Research, Collegeville, PA, 19426, USA  
 SOURCE: Journal of Biological Chemistry (2003), 278(38), 36418-36429  
 CODEN: JBCHA3; ISSN: 0021-9258  
 PUBLISHER: American Society for Biochemistry and Molecular Biology  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Hormonal status can influence diverse metabolic pathways. Small heterodimer partner (SHP) is an orphan nuclear receptor that can modulate the activity of several transcription factors. Estrogens are here shown to directly induce expression of the SHP in the mouse and rat liver and in human HepG2 cells. SHP is rapidly induced within 2 h following treatment of mice with ethynylestradiol (EE) or the estrogen receptor  $\alpha$  (ER $\alpha$ )-selective compound Pr pyrazole triol (PPT). SHP induction by these estrogens is completely absent in ER $\alpha$ KO mice. Mutation of the human SHP promoter defined HNF-3, HNF-4, GATA, and AP-1 sites as important for basal activity, whereas EE induction required two distinct elements located between -309 and -267. One of these elements contains an estrogen response element half-site that bound purified ER $\alpha$ , and ER $\alpha$  with a mutated DNA binding domain was unable to stimulate SHP promoter activity. This ER $\alpha$  binding site overlaps the known **farnesoid X receptor (FXR)** binding site in the SHP promoter, and the combination of EE plus FXR agonists did not produce an additive induction of SHP expression in mice. Surprisingly, induction of SHP by EE did not inhibit expression of the known SHP target genes cholesterol 7 $\alpha$ -hydroxylase (CYP7A1) or sterol 12 $\alpha$ -hydroxylase (CYP8B1). However, the direct regulation of SHP expression may provide a basis for some of the numerous biol. effects of estrogens.  
 IT 278779-30-9, GW4064  
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (estrogen receptor  $\alpha$  regulates expression of orphan receptor small heterodimer partner as studied in mouse and rat liver and in human HepG2 cells)  
 RN 278779-30-9 CAPLUS  
 CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 6 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A


REFERENCE COUNT: 69 THERE ARE 69 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 7 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:698404 CAPLUS  
DOCUMENT NUMBER: 140:87450  
TITLE: Farnesoid X receptor agonists suppress hepatic apolipoprotein CIII expression  
AUTHOR(S): Claudel, Thierry; Inoue, Yumiko; Barbier, Olivier; Duran-Sandoval, Daniel; Kosykh, Vladimir; Fruchart, Jamila; Fruchart, Jean-Charles; Gonzalez, Frank J.; Staels, Bart  
CORPORATE SOURCE: Departement d'Atherosclerose, UR545 INSERM, Institut Pasteur de Lille, Lille, Fr.  
SOURCE: Gastroenterology (2003), 125(2), 544-555  
CODEN: GASTAB; ISSN: 0016-5085  
PUBLISHER: W. B. Saunders Co.  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Background & Aims: Increased serum triglyceride levels constitute a risk factor for coronary heart disease. Apolipoprotein CIII (Apo CIII) is a determinant of serum triglyceride metabolism. In this study, we investigated whether activators of the nuclear farnesoid X receptor (FXR) modulate Apo CIII gene expression. Methods: The influence of bile acids and synthetic FXR activators on Apo CIII and triglyceride metabolism was studied in vivo by using FXR wild-type and FXR-deficient mice and in vitro by using human primary hepatocytes and HepG2 cells. Results: In mice, treatment with the FXR agonist taurocholic acid strongly decreased serum triglyceride levels, an effect associated with reduced Apo CIII serum and liver mRNA levels. By contrast, no change was observed in FXR-deficient mice. Incubation of human primary hepatocytes and HepG2 cells with bile acids or the nonsteroidal synthetic FXR agonist GW4064 resulted in a dose-dependent downregulation of Apo CIII gene expression. Promoter transfection expts. and mutation anal. showed that bile acid-activated FXR decrease human Apo CIII promoter activity via a neg. FXR response element located in the 14 footprint between nucleotides -739 and -704. Chromatin immunoprecipitation expts. showed that bile acid treatment led to binding of FXR/retinoid X receptor heterodimers to and displacement of HNF4α from this site. Bile acid treatment still repressed liver Apo CIII gene expression in hepatic HNF4α-deficient mice, suggesting an active rather than a competitive mechanism of Apo CIII repression by the FXR. Conclusions: We identified bile acid and synthetic activators of the nuclear FXR as neg. regulators of Apo CIII expression, an effect that may contribute to the triglyceride-decreasing action of FXR agonists.  
IT 278779-30-9, GW4064  
RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(farnesoid X receptor agonists suppress hepatic apolipoprotein CIII expression)  
RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 7 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A


REFERENCE COUNT: 71 THERE ARE 71 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 8 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:579493 CAPLUS  
DOCUMENT NUMBER: 139:256039  
TITLE: Human kininogen gene is transactivated by the farnesoid X receptor  
AUTHOR(S): Zhao, Annie; Lew, Jane-L.; Huang, Li; Yu, Jinghua; Zhang, Theresa; Hrymka, Yaroslava; Thompson, John R.; de Pedro, Nuria; Blevins, Richard A.; Pelaez, Fernando; Wright, Samuel D.; Cui, Jisong  
CORPORATE SOURCE: Departments of Atherosclerosis and Endocrinology, Merck Research Laboratories, Rahway, NJ, 07065, USA  
SOURCE: Journal of Biological Chemistry (2003), 278(31), 28765-28770  
CODEN: JBCHA3; ISSN: 0021-9258  
PUBLISHER: American Society for Biochemistry and Molecular Biology  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Human kininogen belongs to the plasma kallikrein-kinin system. High mol. weight kininogen is the precursor for two-chain kinin-free kininogen and bradykinin. It has been shown that the two-chain kinin-free kininogen has the properties of anti-adhesion, anti-platelet aggregation, and anti-thrombosis, whereas bradykinin is a potent vasodilator and mediator of inflammation. In this study the human kininogen gene is strongly up-regulated by agonists of the farnesoid X receptor (FXR), a nuclear receptor for bile acids. In primary human hepatocytes, both the endogenous FXR agonist chenodeoxycholate and synthetic FXR agonist GW4064 increased kininogen mRNA with a maximum induction of 8-10-fold. A more robust induction of kininogen expression was observed in HepG2 cells, where kininogen mRNA was increased by chenodeoxycholate or GW4064 up to 130-140-fold as shown by real time PCR. Northern blot anal. confirmed the up-regulation of kininogen expression by FXR agonists. To determine whether kininogen is a direct target of FXR, the authors examined the sequence of the kininogen promoter and identified a highly conserved FXR response element (inverted repeat, IR-1) in the proximity of the kininogen promoter (-66/-54). FXR/RXRα heterodimers specifically bind to this IR-1. A construct of a minimal promoter with the luciferase reporter containing this IR-1 was transactivated by FXR. Deletion or mutation of this IR-1 abolished FXR-mediated promoter activation, indicating that this IR-1 element is responsible for the promoter transactivation by FXR. The authors conclude that kininogen is a novel and direct target of FXR, and bile acids may play a role in the vasodilation and anti-coagulation processes.  
IT 278779-30-9, GW4064  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (human kininogen gene is transactivated by the farnesoid X receptor in primary human hepatocytes)  
RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 8 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A

REFERENCE COUNT:


45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 9 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:375244 CAPLUS  
DOCUMENT NUMBER: 139:159454  
TITLE: A chemical, genetic, and structural analysis of the nuclear bile acid receptor FXR  
AUTHOR(S): Downes, Michael; Verdecia, Mark A.; Roecker, A. J.; Hughes, Robert; Hogenesch, John B.; Kast-Woelbern, Heidi R.; Bowman, Marianne E.; Ferrer, Jean-Luc; Anisfeld, Andrew M.; Edwards, Peter A.; Rosenfeld, John M.; Alvarez, Jacqueline G. A.; Noel, Joseph P.; Nicolaou, K. C.; Evans, Ronald M.  
CORPORATE SOURCE: Gene Expression Laboratory, Howard Hughes Medical Institute, La Jolla, CA, 92037, USA  
SOURCE: Molecular Cell (2003), 11(4), 1079-1092  
CODEN: MOCEFL; ISSN: 1097-2765  
PUBLISHER: Cell Press  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The farnesoid X receptor (FXR) functions as a bile acid (BA) sensor coordinating cholesterol metabolism, lipid homeostasis, and absorption of dietary fats and vitamins. However, BAs are poor reagents for characterizing FXR functions due to multiple receptor independent properties. Accordingly, using combinatorial chemical we evolved a small mol. agonist termed fexaramine with 100-fold increased affinity relative to natural compds. Gene-profiling expts. conducted in hepatocytes with FXR-specific fexaramine vs. the primary BA chenodeoxycholic acid (CDCA) produced remarkably distinct genomic targets. Highly diffracting cocrystals (1.78 Å) of fexaramine bound to the ligand binding domain of FXR revealed the agonist sequestered in a 726 Å<sup>3</sup> hydrophobic cavity and suggest a mechanistic basis for the initial step in the BA signaling pathway. The discovery of fexaramine will allow us to unravel the FXR genetic network from the BA network and selectively manipulate components of the cholesterol pathway that may be useful in treating cholesterol-related human diseases.  
IT 278779-30-9, GW 4064  
RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (chemical, genetic, and structural anal. of nuclear bile acid receptor FXR)  
RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 9 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A

REFERENCE COUNT:


39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 10 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:237176 CAPLUS  
DOCUMENT NUMBER: 139:17879  
TITLE: Differential regulation of rat and human CYP7A1 by the nuclear oxysterol receptor liver X receptor-α  
AUTHOR(S): Goodwin, Bryan; Watson, Michael A.; Kim, Hwajin; Miao, Ji; Kemper, Jongsook Kim; Kliewer, Steven A.  
CORPORATE SOURCE: Nuclear Receptor Discovery Research, GlaxoSmithKline Research and Development, Research Triangle Park, NC, 27709, USA  
SOURCE: Molecular Endocrinology (2003), 17(3), 386-394  
CODEN: MOENEN; ISSN: 0888-8809  
PUBLISHER: Endocrine Society  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB In rodent liver, transcription of the gene encoding cholesterol 7α-hydroxylase (CYP7A1), which catalyzes the rate-limiting step in the classic bile acid synthetic pathway, is stimulated by the liver X receptor α (LXRα), a nuclear receptor for oxysterol metabolites of cholesterol. This feed-forward regulatory loop provides a mechanism for the elimination of excess cholesterol from the body. The authors demonstrate that in primary cultures of human hepatocytes, activation of LXRα has the opposite effect, repressing CYP7A1 expression. This repression is mediated, at least in part, through induction of the orphan nuclear receptor, short heterodimer partner (SHP), which is also induced by bile acids. The authors demonstrate that SHP is regulated directly by LXRα through a DNA response element that overlaps with the previously characterized bile acid response element. The authors' data reveal a fundamental difference in the regulation of CYP7A1 in rodent and human hepatocytes and provide evidence that different species employ distinct mol. strategies to regulate cholesterol homeostasis.  
IT 278779-30-9, GW4064  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (differential regulation of rat and human CYP7A1 by nuclear oxysterol receptor liver X receptor-α)  
RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 10 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A

REFERENCE COUNT:

49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 11 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:204786 CAPLUS

DOCUMENT NUMBER: 139:79298

TITLE: Guggulsterone Is a Farnesoid X Receptor Antagonist in Coactivator Association Assays but Acts to Enhance Transcription of Bile Salt Export Pump

AUTHOR(S): Cui, Jisong; Huang, Li; Zhao, Annie; Lew, Jane-L.; Yu, Jinghua; Sahoo, Soumya; Meinke, Peter T.; Royo, Inmaculada; Pelaez, Fernando; Wright, Samuel D.

CORPORATE SOURCE: Department of Atherosclerosis and Endocrinology, Merck Research Laboratories, Rahway, NJ, 07065, USA

SOURCE: Journal of Biological Chemistry (2003), 278(12), 10214-10220

CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER: American Society for Biochemistry and Molecular Biology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Guggulipid is an extract of the guggul tree *Commiphora mukul* and has been widely used to treat hyperlipidemia in humans. The plant sterol guggulsterone (GS) is the active agent in this extract. Recent studies have shown that GS can act as an antagonist ligand for farnesoid X receptor (FXR) and decrease expression of bile acid-activated genes. Here we show that GS, although an FXR antagonist in coactivator association

assays, enhances FXR agonist-induced transcription of bile salt export pump (BSEP), a major hepatic bile acid transporter. In HepG2 cells, in the presence of an FXR agonist such as chenodeoxycholate or GW4064, GS enhanced endogenous BSEP expression with a maximum induction of 400-500% that


induced by an FXR agonist alone. This enhancement was also readily observed in FXR-dependent BSEP promoter activation using a luciferase reporter construct. In addition, GS alone slightly increased BSEP promoter activation in the absence of an FXR agonist. Consistent with the results in HepG2, guggulipid treatment in Fisher rats increased BSEP mRNA. Interestingly, in these animals expression of the orphan nuclear receptor SHP (small heterodimer partner), a known FXR target, was also significantly increased, whereas expression of other FXR targets including cholesterol 7 $\alpha$ -hydroxylase (Cyp 7a1), sterol 12 $\alpha$ -hydroxylase (Cyp 8b1), and the intestinal bile acid-binding protein (I-BAP), remained unchanged. Thus, we propose that GS is a selective bile acid receptor modulator that regulates expression of a subset of FXR targets. Guggulipid treatment in rats lowered serum triglyceride and raised serum high d. lipoprotein levels. Taken together, these data suggest that guggulsterone defines a novel class of FXR ligands characterized by antagonist activities in coactivator association assays but with the ability to enhance the action of agonists on BSEP expression in vivo.

IT 278779-30-9, GW4064  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (FXR agonist: guggulsterone is a farnesoid X receptor antagonist in coactivator association assays but Acts to enhance transcription of bile salt export pump)

RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[(3-[2,6-dichlorophenyl]-5-(1-methylethyl)-4-isoxazolyl)methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 11 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A

REFERENCE COUNT:

44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 12 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:112477 CAPLUS

DOCUMENT NUMBER: 138:298694

TITLE: Bile acids induce the expression of the human peroxisome proliferator-activated receptor  $\alpha$  gene via activation of the farnesoid X receptor

AUTHOR(S): Torta, Ines Pineda; Claudel, Thierry; Duval, Caroline; Kosykh, Vladimir; Fruchart, Jean-Charles; Staels, Bart

CORPORATE SOURCE: U.545 Institut National de la Sante et de la Recherche Medicale, Departement d'Atherosclerosis, Institut Pasteur de Lille, Lille, 59019, Fr.

SOURCE: Molecular Endocrinology (2003), 17(2), 259-272

CODEN: MOENEN; ISSN: 0898-8809

PUBLISHER: Endocrine Society

DOCUMENT TYPE: Journal

LANGUAGE: English


AB Peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ) is a nuclear receptor that controls lipid and glucose metabolism and exerts antiinflammatory activities. PPAR $\alpha$  is also reported to influence bile acid formation and bile composition. Farnesoid X receptor (FXR) is a bile acid-activated nuclear receptor that mediates the effects of bile acids on gene expression and plays a major role in bile acid and possibly also in lipid metabolism. Thus, both PPAR $\alpha$  and FXR appear to act on common metabolic pathways. To determine the existence of a mol. cross-talk between these two nuclear receptors, the regulation of PPAR $\alpha$  expression by bile acids was investigated. Incubation of human hepatoma HepG2 cells with the natural FXR ligand chenodeoxycholic acid (CDCA) as well as with the nonsteroidal FXR agonist GW4064 resulted in a significant induction of PPAR $\alpha$  mRNA levels. In addition, hPPAR $\alpha$  gene expression was up-regulated by taurocholic acid in human primary hepatocytes. Cotransfection of FXR/retinoid X receptor in the presence of CDCA led to up to a 3-fold induction of human PPAR $\alpha$  promoter activity in HepG2 cells. Mutation anal. identified a FXR response element ( $\alpha$ FXRE) that mediates bile acid regulation of this promoter. FXR bound the  $\alpha$ FXRE site as demonstrated by gel shift anal., and CDCA specifically increased the activity of a heterologous promoter driven by four copies of the  $\alpha$ FXRE. In contrast, neither the murine PPAR $\alpha$  promoter, in which the  $\alpha$ FXRE is not conserved, nor a mouse  $\alpha$ FXRE-driven heterologous reporter, were responsive to CDCA treatment. Moreover, PPAR $\alpha$  expression was not regulated in taurocholic acid-fed mice. Finally, induction of hPPAR $\alpha$  mRNA levels by CDCA resulted in an enhanced induction of the expression of the PPAR $\alpha$  target gene carnitine palmitoyltransferase I by PPAR $\alpha$  ligands. In concert, these results demonstrate that bile acids stimulate PPAR $\alpha$  expression in a species-specific manner via a FXRE located within the human PPAR $\alpha$  promoter. These results provide mol. evidence for a cross-talk between the FXR and PPAR $\alpha$  pathways in humans.

IT 278779-30-9, GW4064  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses) (PPAR $\alpha$  mRNA induction by: bile acids induce the expression of the human peroxisome proliferator-activated receptor  $\alpha$  gene via activation of the farnesoid X receptor)

RN 278779-30-9 CAPLUS  
CN Benzoic acid, 3-[2-[2-chloro-4-[(3-[2,6-dichlorophenyl]-5-(1-methylethyl)-4-isoxazolyl)methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 12 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A

REFERENCE COUNT: 69 THERE ARE 69 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 13 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:677926 CAPLUS

DOCUMENT NUMBER: 138:49877

TITLE: Lithocholic acid decreases expression of bile salt export pump through farnesoid X receptor antagonist activity

AUTHOR(S): Yu, Jinghua; Lo, Jane-L.; Huang, Li; Zhao, Annie; Metzger, Edward; Adams, Alan; Meinke, Peter T.; Wright, Samuel D.; Cui, Jisong

CORPORATE SOURCE: Department of Atherosclerosis and Endocrinology, Merck Research Laboratories, Rahway, NJ, 07065, USA

SOURCE: Journal of Biological Chemistry (2002), 277(35), 31441-31447

CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER: American Society for Biochemistry and Molecular Biology

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Bile salt export pump (BSEP) is a major bile acid transporter in the liver. Mutations in BSEP result in progressive intrahepatic cholestasis, a severe liver disease that impairs bile flow and causes irreversible liver damage. BSEP is a target for inhibition and down-regulation by drugs and abnormal bile salt metabolites, and such inhibition and down-regulation may result in bile acid retention and intrahepatic cholestasis. In this study, we quant. analyzed the regulation of BSEP expression by FXR ligands in primary human hepatocytes and HepG2 cells. We demonstrate that BSEP expression is dramatically regulated by ligands of the nuclear receptor farnesoid X receptor (FXR). Both the endogenous FXR agonist chenodeoxycholate (CDCA) and synthetic FXR ligand GW4064 effectively increased BSEP mRNA in both cell types. This up-regulation was readily detectable at as early as 3 h, and the ligand potency for BSEP regulation correlates with the intrinsic activity on FXR. These results suggest BSEP as a direct target of FXR and support the recent report that the BSEP promoter is transactivated by FXR. In contrast to CDCA and GW4064, lithocholate (LCA), a hydrophobic bile acid and a potent inducer of cholestasis, strongly decreased BSEP expression. Previous studies did not identify LCA as an FXR antagonist ligand in cells, but we show here that LCA is an FXR antagonist with partial agonist activity in cells. In an in vitro coactivator association assay, LCA decreased CDCA- and GW4064-induced FXR activation with an IC50 of 1 μM. In HepG2 cells, LCA also effectively antagonized GW4064-enhanced FXR transactivation. These data suggest that the toxic and cholestatic effect of LCA in animals may result from its down-regulation of BSEP through FXR. Taken together, these observations indicate that FXR plays an important role in BSEP gene expression and that FXR ligands may be potential therapeutic drugs for intrahepatic cholestasis.

IT 278779-30-9, GW4064


RL: BSU (Biological study, unclassified); BIOL (Biological study) (endogenous FXR agonist chenodeoxycholate and synthetic FXR ligand GW4064 effectively increases BSEP (bile salt export pump) mRNA in primary human hepatocytes and HepG2 cells)

RN 278779-30-9 CAPLUS

CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 13 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A

REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 14 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2002:540315 CAPLUS

DOCUMENT NUMBER: 137:263227

TITLE: 6α-Ethyl-Chenodeoxycholic Acid (6-ECDCA), a Potent and Selective FXR Agonist Endowed with Anticholestatic Activity

AUTHOR(S): Pallicciari, Roberto; Fiorucci, Stefano; Cernaioni, Emidio; Clerici, Carlo; Costantino, Gabriele; Maloney, Patrick R.; Morelli, Antonio; Parks, Derek J.; Willson, Timothy M.

CORPORATE SOURCE: Dipartimento di Chimica e Tecnologia del Farmaco, Università di Perugia, Perugia, 06123, Italy

SOURCE: Journal of Medicinal Chemistry (2002), 45(17), 3569-3572


CODEN: JMCMAH; ISSN: 0022-2623

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



AB A series of 6α-alkyl-substituted analogs I (R = Me, Et, Pr, Bu) of chenodeoxycholic acid (CDCA) were synthesized and evaluated as potential farnesoid X receptor (FXR) ligands. Among them, 6α-ethyl-chenodeoxycholic acid (6-ECDCA) I (R = Et) was shown to be a very potent and selective FXR agonist (EC50 = 99 nM) and to be endowed with anticholestatic activity in an in vivo rat model of cholestasis.

IT 278779-30-9, GW4064


RL: BSU (Biological study, unclassified); BIOL (Biological study) (GW 4064; binding potency to farnesoid X receptor agonist endowed with anticholestatic activity)

RN 278779-30-9 CAPLUS

CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 14 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A

REFERENCE COUNT:

17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 15 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2001:729132 CAPLUS

DOCUMENT NUMBER: 136:18310

TITLE: Farnesoid X-activated receptor induces apolipoprotein C-II transcription: a molecular mechanism linking plasma triglyceride levels to bile acids

AUTHOR(S): Kast, Heidi Rachelle; Nguyen, Catherine M.; Sinal, Christopher J.; Jones, Stacey A.; Laffitte, Bryan A.; Reue, Karen; Gonzalez, Frank J.; Willson, Timothy M.; Edwards, Peter A.

CORPORATE SOURCE: Departments of Biological Chemistry and Medicine, University of California, Los Angeles, CA, 90095, USA

SOURCE: Molecular Endocrinology (2001), 15(10), 1720-1728

CODEN: MOENEN; ISSN: 0898-8809

PUBLISHER: Endocrine Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The farnesoid X-activated receptor (FXR; NR1H4), a member of the nuclear hormone receptor superfamily, induces gene expression in response to several bile acids, including chenodeoxycholic acid. Here the authors used suppression subtractive hybridization to identify apolipoprotein C-II (apoC-II) as an FXR target gene. Retroviral expression of FXR in HepG2 cells results in induction of the mRNA encoding apoC-II in response to several FXR ligands. EMSAs demonstrate that recombinant FXR and RXR bind to two FXR response elements that are contained within two important distal enhancer elements (hepatic control regions) that lie 11 kb and 22 kb upstream of the transcription start site of the apoC-II gene. A luciferase reporter gene containing the hepatic control region or two copies of the wild-type FXR response element was activated when FXR-containing

cells were treated with FXR ligands. In addition, the authors report that hepatic expression of both apoC-II and phospholipid transfer protein mRNAs increases when mice are fed diets supplemented with cholic acid, an FXR ligand, and this induction is attenuated in FXR null mice. Finally, the authors observed decreased plasma triglyceride levels in mice fed cholic acid-containing diets. These results identify a mechanism whereby FXR and its ligands lower plasma triglyceride levels. These findings may have important implications in the clin. management of hyperlipidemias.

IT 278779-30-9, GW 4064

RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); BIOL (Biological study)


(farnesoid X-activated receptor induces apolipoprotein C-II transcription in HepG2 cells in relation to mol. mechanism linking plasma triglyceride levels to bile acids)

RN 278779-30-9 CAPLUS

CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)

L7 ANSWER 15 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

PAGE 1-A



PAGE 2-A

REFERENCE COUNT:

36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 16 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2000:441628 CAPLUS

DOCUMENT NUMBER: 133:68969

TITLE: Assays for ligands for nuclear receptors using peptide sequences

INVENTOR(S): Blanchard, Steven Gerard; Klievar, Anthony; Lehmann, Jurgen; Parks, Derek J.; Stimmel, Julie Beth; Willson, Timothy Mark

PATENT ASSIGNEE(S): Glaxo Group Limited, UK

SOURCE: PCT Int. Appl., 62 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

| PATENT NO.  | KIND | DATE     | APPLICATION NO. | DATE     |
|---|------|----------|-----------------|----------|
| WO 2000037077   | A1   | 20000629 | WO 1999-US30947 | 19991222 |
| W: AE, AL, AM, AT, AU, AZ, BG, BR, CA, CH, CN, CU, DE, DK, EE, ES, FI, GB, GD, GH, HR, IN, IS, JP, KE, KU, LV, MD, MN, MW, MX, NO, RU, SD, SE |      |          |                 |          |
| RW: GH, GM, KE, LS, MW, SD, SL, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, HR, NE, TD, TG  |      |          |                 |          |
| CA 2356887  | AA   | 20000629 | CA 1999-2356887 | 19991222 |
| EP 1140079  | A1   | 20011010 | EP 1999-967639  | 19991222 |
| R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO   |      |          |                 |          |
| JP 2002532729   | T2   | 20021002 | JP 2000-589188  | 19991222 |
| US 6639078  | B1   | 20031028 | US 2001-868397  | 20010618 |
| US 2004048316   | A1   | 20040311 | US 2003-637190  | 20030808 |
| PRIORITY APPLN. INFO.:  |      |          |                 |          |
| US 1998-135097P P 19981223  |      |          |                 |          |
| WO 1999-US30947 W 19991222  |      |          |                 |          |
| US 2001-868397 A1 20010618  |      |          |                 |          |

OTHER SOURCE(S): MARPAT 133:68969


AB The present invention provides a method of identifying compds. for the treatment of diseases or disorders modulated by farnesoid X receptor (FXR), comprising the step of determining whether the compound interacts directly with FXR, wherein a compound that interacts directly with FXR is a compound for the treatment. A generic approach to assay development for nuclear receptors is presented, using purified ligand binding domains. The concept of generic assay development is extended to develop in vitro assays that detect ligand binding by monitoring ligand-induced changes in receptor heterodimerization. This approach is demonstrated using both scintillation proximity and homogeneous time-resolved fluorimetry (HTRF). Another aspect of the invention is a nuclear receptor peptide assay for identifying ligands. This assay utilizes fluorescence resonance energy transfer (FRET) and can be used to test whether putative ligands bind to FXR. The FRET assay is based upon the principle that ligands induce conformational changes in nuclear receptors that facilitate interactions with coactivator proteins required for transcriptional activation. Binding of the FXR nuclear receptor can result in the alteration of expression of various genes that FXR aids in regulating, including genes involved in lipid absorption and digestion in the small intestine and lipid homeostasis in liver. FXR often functions as a heterodimer with the RXR receptor. The inventive method includes using this technol. to affect bile acid and cholesterol homeostasis such that, ultimately, cholesterol and lipid levels can be modified and in treating diseases in a mammal, including human, in which regulation of bile acid, cholesterol and lipid levels is important. For example, GW4064 (prepared in a yield of 98%) was



L7 ANSWER 16 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)  
 given to Fischer rats at a dose of 30 mg/kg for 7 days. At the end of  
 study, serum triglyceride levels were decreased by 264 compared to a  
 vehicle-treated controls. Nearly 20 genes were identified in the  
 intestine that were regulated 51.5-fold by GW4064. The expression of  
 roughly half of these genes was decreased by GW4064 treatment. All of  
 these down-regulated genes are involved in either lipid absorption or  
 proteolysis, including lipases, proteases, and a colipase.

IT 278779-30-9P, GW 4064  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological  
 process); BSU (Biological study, unclassified); SPN (Synthetic  
 preparation); THU (Therapeutic use); BIOL (Biological study); PREP  
 (Preparation); PROC (Process); USES (Uses)  
 (identification of nuclear receptor ligands for treatment of diseases  
 affected by cholesterol, triglycerides and bile acid levels)

RN 278779-30-9 CAPLUS  
 CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-  
 4-isoxazolyl]methoxy]phenyl]ethenyl]- (9CI) (CA INDEX NAME)



PAGE 1-A




PAGE 2-A

IT 278779-31-0P, GW 4064 methyl ester  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)

L7 ANSWER 16 OF 16 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)  
 (prepn. of GW4064 as nuclear ~~l~~arnesoid X receptor ligand)

RN 278779-31-0 CAPLUS  
 CN Benzoic acid, 3-[2-[2-chloro-4-[[3-(2,6-dichlorophenyl)-5-(1-methylethyl)-  
 4-isoxazolyl]methoxy]phenyl]ethenyl]-, methyl ester (9CI) (CA INDEX NAME)



PAGE 1-A



PAGE 2-A

REFERENCE COUNT:

4

THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

78.80

251.07

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-11.09

-11.09

STN INTERNATIONAL LOGOFF AT 13:02:15 ON 23 JUN 2004